

# Skew braoids and Hopf-Galois structures

Paul Truman

Keele University, UK

18th April, 2023

# Overview

- Joint work with Isabel Martin-Lyons


## Aim

Generalize the definition of skew braces to give objects corresponding to Hopf-Galois structures on separable, but non-normal, extensions.

- A route for constructing a skew brace from a Hopf-Galois structure on a Galois extension.
- Generalizing to skew bracoids
- Characterizations and  $\gamma$ -functions
- Ideals and quotients
- Hopf-Galois structures and the Hopf-Galois correspondence

## Greither-Pareigis theory for non-normal extensions

- Let  $L/K$  be a separable extension of fields with Galois closure  $E$ .
- Write  $G = \text{Gal}(E/K)$  and  $G_L = \text{Gal}(E/L)$ .
- Let  $X = G/G_L$  and define  $\lambda : G \rightarrow \text{Perm}(X)$  by  $\lambda(g)[\bar{h}] = \overline{gh}$ .
- Then  $G$  acts on  $\text{Perm}(X)$  by conjugation via  $\lambda$ .
- There is a bijection between  $G$ -stable regular subgroups of  $\text{Perm}(X)$  and Hopf-Galois structures on  $L/K$ .



## The S-T route from a HGS to a skew brace

- Let  $L/K$  be a Galois extension with Galois group  $G = (G, \cdot)$ .
- Suppose that  $N$  is a regular  $G$ -stable subgroup of  $\text{Perm}(G)$ .
- The map  $N \rightarrow G$  defined by  $\eta \mapsto \eta[e_G]$  is a bijection.
- Transport the structure of  $N^{opp}$  to  $G$  via

$$\eta[e_G] \star \mu[e_G] = (\mu\eta)[e_G].$$

- Then  $(G, \star)$  is a group isomorphic to  $N$  and

$$g \cdot (h_1 \star h_2) = (g \cdot h_1) \star g^{-1} \star (g \cdot h_2),$$

so  $(G, \star, \cdot)$  is a skew brace.

## Mimicking the route in the non-normal case

- Now let  $L/K$  be separable, but non-normal, with Galois closure  $E$ .
- Suppose that  $N$  is a regular  $G$ -stable subgroup of  $\text{Perm}(X)$ .
- The map  $N \rightarrow X$  defined by  $\eta \mapsto \eta[\overline{e}_G]$  is a bijection.
- Transport the structure of  $N^{opp}$  to  $X$  via

$$\eta[\overline{e}_G] \star \mu[\overline{e}_G] = (\mu\eta)[\overline{e}_G].$$

- Then  $(X, \star)$  is a group isomorphic to  $N$  and

$$g \odot (\overline{x}_1 \star \overline{x}_2) = (g \odot \overline{x}_1) \star \overline{g}^{-1} \star (g \odot \overline{x}_2),$$

where  $\odot$  denotes left translation of cosets.

# Skew bracoids

## Definition

A *skew bracoid* is a 5-tuple  $(G, \cdot, N, \star, \odot)$  where  $(G, \cdot)$  and  $(N, \star)$  are groups and  $\odot$  is a transitive action of  $(G, \cdot)$  on  $N$  such that

$$g \odot (\eta \star \mu) = (g \odot \eta) \star (g \odot e_N)^{-1} \star (g \odot \mu) \quad (*)$$

for all  $g \in G$  and  $\eta, \mu \in N$ .

- Where possible, we write simply  $(G, N, \odot)$ , or even  $(G, N)$ .
- For now, we always assume  $G, N$  are finite. Then  $|G| = |S||N|$ , where  $S = \text{Stab}_G(e_N)$ .
- Every skew brace is a skew bracoid, with  $\odot$  and  $\cdot$  coinciding.
- If  $|N| = |G|$  then  $(G, N)$  is essentially a skew brace.

## An example

### Example

- Let  $(G, \star, \cdot)$  be a skew brace and let  $H$  be a strong left ideal.
- $H$  is a normal subgroup of  $(G, \star)$ , so  $(G/H, \star)$  is a group.
- $H$  is a subgroup of  $(G, \cdot)$ , so  $(G, \cdot)$  acts by left translation on the coset space  $G/H$ . Write  $\odot$  for this action.
- Then  $(G, \cdot, G/H, \star, \odot)$  is a skew bracoid.

### Question

Does every skew bracoid occur in this way?

## Some characterizations

### Proposition

Let  $(G, \cdot)$  and  $(N, \star)$  be groups, let  $\odot$  be a transitive action of  $G$  on  $N$ , and let

$$\lambda_{\odot} : G \rightarrow \text{Perm}(N)$$

be the corresponding permutation representation. Then  $(G, N)$  is a skew bracoid if and only if  $\lambda_{\odot}(G) \subseteq \text{Hol}(N)$ .

### Definition

A skew bracoid  $(G, N)$  is called *reduced* if  $\lambda_{\odot} : G \rightarrow \text{Hol}(N)$  is injective.

### Proposition

Let  $(N, \star)$  be a group. There is a bijective correspondence between transitive subgroups  $G$  of  $\text{Hol}(N)$  and reduced skew bracoids  $(G, N)$ .



# Equivalence

## Definition

Two skew bracoids  $(G, N)$  and  $(G', N')$  are called *equivalent* if  $N = N'$  and  $\lambda_{\odot}(G) = \lambda_{\odot'}(G') \subseteq \text{Hol}(N)$ .

## Proposition

Let  $(G, N)$  be a skew bracoid.

- Let  $\overline{G} = G / \ker(\lambda_{\odot})$ . Then  $(\overline{G}, N)$  is a reduced skew bracoid, which will be called the *reduced form* of  $(G, N)$ .
- Every skew bracoid is equivalent to its reduced form.

## $\gamma$ -functions

### Proposition

Let  $(G, N)$  be a skew braceoid. Define  $\gamma : G \rightarrow \text{Perm}(N)$  by

$$\gamma(g)\eta = (g \odot e_N)^{-1} \star (g \odot \eta) \text{ for all } g \in G \text{ and } \eta \in N.$$

Then

- $\gamma$  is a group homomorphism;
  - $\gamma(G) \subseteq \text{Aut}(N)$ .
- 
- Caranti uses  $\gamma : G \rightarrow \text{Perm}(G)$  to characterize regular subgroups of  $\text{Hol}(G)$ , and hence skew braces. This approach does not seem to work well for transitive subgroups and skew braceoids.
  - But see Stefanelli: affine structures etc.

# Substructures

## Definition

A *sub skew bracoid* of a skew bracoid  $(G, N)$  consists of a subgroup  $H$  of  $G$  and a subgroup  $M$  of  $N$  such that  $(H, M)$  is a skew bracoid.

- It is possible for  $(G, N)$  to be reduced but for  $(H, M)$  to not be so.

## Definition

A *left ideal* of a skew bracoid  $(G, N)$  is a subgroup  $M$  of  $N$  such that  $\gamma^{(g)}M = M$  for all  $g \in G$ . An *ideal* is a left ideal  $M$  that is normal in  $N$ .

## Proposition

If  $M$  is an ideal of  $(G, N)$  then  $(G, N/M)$  is a skew bracoid.

# Ideals

## Proposition

Let  $M$  be a left ideal of  $(G, N)$ , and let

$$G_M = \{g \in G \mid g \odot \mu \in M \text{ for all } \mu \in M\}.$$

Then  $(G_M, M)$  is a sub skew bracoid of  $(G, N)$ .

## Proof.

It is clear that  $G_M$  is a subgroup of  $G$ .

Let  $G_M^{(e)} = \{g \in G \mid g \odot e_N \in M\}$ .

For all  $g \in G$  and  $\mu \in M$  we have

$$\gamma(g)\mu = (g \odot e_N)^{-1} \star (g \odot \mu) \in M.$$

Hence  $G_M^{(e)} = G_M$ , so  $G_M$  is transitive on  $M$ . □

## Back to Hopf-Galois structures

If  $L/K$  is a Galois extension with Galois group  $(G, \cdot)$ , Stefanello and Trappeniers show that there is a bijection between

- binary operations  $\star$  on  $G$  such that  $(G, \star, \cdot)$  is a skew brace;
- Hopf-Galois structures on  $L/K$ ,

and also that the Hopf-Galois structure corresponding to  $(G, \star, \cdot)$  is given by  $L[G, \star]^{\gamma(G)}$ , acting via

$$\left( \sum_{g \in G} c_g g \right) [t] = \sum_{g \in G} c_g g[t].$$

## Back to Hopf-Galois structures


### Theorem

Let  $E/K$  be a Galois extension with Galois group  $G$ , let  $S \leq G$ . There is a bijection between

- binary operations  $\star$  on  $X = G/S$  such that  $(G, \cdot, X, \star, \odot)$  is a skew bracoid;
- Hopf-Galois structures on  $E^S/K$ .

### Proof.

We have already seen how to get from HGS on  $E^S/K$  to skew bracoid  $(G, \cdot, X, \star, \odot)$ .



## Back to Hopf-Galois structures

### Proof continued...

Conversely, given a skew bracoid of the form  $(G, \cdot, X, \star, \odot)$ , consider

$\rho_\star : X \rightarrow \text{Perm}(X)$  defined by  $\rho_\star(\bar{x})[\bar{y}] = \bar{y} \star \bar{x}^{-1}$ .

Then  $\rho_\star(X)$  is a regular subgroup of  $\text{Perm}(X)$ , and

$$\begin{aligned}\lambda_\odot(g)\rho_\star(\bar{x})\lambda_\odot(g^{-1})[\bar{y}] &= g \odot ((g^{-1} \odot \bar{y}) \star \bar{x}^{-1}) \\ &= \bar{y} \star (g \odot \bar{e})^{-1} \star (g \odot \bar{x}^{-1}) \\ &= \bar{y} \star ((g \odot \bar{e})^{-1} \star (g \odot \bar{x})^{-1}) \\ &= \rho_\star((g \odot \bar{e})^{-1} \star (g \odot \bar{x}))[\bar{y}] \\ &= \rho_\star\left(\gamma^{(g)}\bar{x}\right)[\bar{y}]\end{aligned}$$

So  $\rho_\star(X)$  is  $G$ -stable, and therefore corresponds to a HGS on  $E^S/K$ .  $\square$

## Hopf algebras and subalgebras

If  $L/K$  is a Galois extension with Galois group  $(G, \cdot)$  and  $(G, \star, \cdot)$  is a skew brace, Stefanello and Trappeniers show that

- The intermediate fields realized by the HGS  $L[G, \star]^{\gamma(G)}$  correspond with sub Hopf algebras, which correspond with left ideals of  $(G, \star, \cdot)$ .
- If  $(G', \star, \cdot)$  is a left ideal of  $(G, \star, \cdot)$  then  $L^{(G', \star)} = L^{(G', \cdot)}$ .
- We obtain a quotient HGS on  $L^{G'}/K$  if and only if  $(G', \star, \cdot)$  is a strong left ideal, and a quotient skew brace if and only if  $(G', \star, \cdot)$  is an ideal.



## Hopf algebras and subalgebras

Now let  $E/K$  be Galois with group  $G$ , let  $S \leq G$ , let  $L = E^S$ , and let  $(G, \cdot, X, \star, \odot)$  be a skew bracoid

### Theorem

- *The corresponding Hopf-Galois structure on  $E^S/K$  is given by  $E[X, \star]^{\gamma(G)}$ , acting via*

$$\left( \sum_{\bar{x} \in X} c_{\bar{x}} \bar{x} \right) [t] = \sum_{\bar{x} \in X} c_{\bar{x}} \bar{x}[t].$$

- *continued...*

# Hopf algebras and subalgebras

## Theorem ( continued...)

- *The intermediate fields realized by the HGS correspond with left ideals of  $(G, \cdot, X, \star, \odot)$ .*
- *The left ideals have the form  $X' = G'/S$  for certain  $G' \leq G$ , and  $L(X', \star) = E(G', \cdot)$ .*
- *We obtain a quotient HGS on  $L^{G'}/K$  and a quotient skew bracoid if and only if  $X'$  is an ideal.*
- *Of course, the extension  $L^{G'}/K$  is Galois if and only if  $G'$  is normal in  $G$ .*

# The natural question

## Question

Do skew bracoids have anything to do with the Yang-Baxter equation?

- I'm working on it!

Thank you for your attention.